

Equidimensional Symmetric Algebras
and Residual Intersections

by

Mark R. Johnson

UofA-R-188

Equidimensional Symmetric Algebras and Residual Intersections

Mark R. Johnson

Abstract. For a finitely generated module M , over a universally catenary local ring, whose symmetric algebra is equidimensional, the ideals generated by the rows of a minimal presentation matrix are shown to have height at most $\mu(M) - \text{rank } M$. Moreover, in the extremal case, they are Cohen-Macaulay ideals if the symmetric algebra is Cohen-Macaulay. Some applications are given to residual intersections of ideals.

Let R be a noetherian ring and let M be a finitely generated R -module. If

$$R^m \xrightarrow{\phi} R^n \rightarrow M \rightarrow 0$$

is a presentation of M , then one may define the symmetric algebra $S(M)$ of M as

$$S(M) = R[T_1, \dots, T_n] / (\ell_1, \dots, \ell_m),$$

where

$$[\ell_1, \dots, \ell_m] = [T_1, \dots, T_n] \cdot \phi.$$

Thus the properties of $S(M)$ are reflected in the presentation matrix ϕ of M .

The problem of determining when the symmetric algebra is a domain has been extensively studied over the years. Although a great deal is known, one does not have a definitive answer. On the other hand, quite generally one knows that if $\text{Spec}(S(M))$ is irreducible then $S(M)$ is equidimensional ([13], 2.1.1). In this paper, we consider the latter condition and the constraint it places on the matrix ϕ . Our main result is the following, which we state in the local case.

THEOREM 1. Let R be a universally catenary local ring, let M be a finitely generated R -module with presentation $R^m \xrightarrow{\phi} R^n \rightarrow M \rightarrow 0$, and let $J \neq R$ be the ideal generated by a row of ϕ . If $S(M)$ is equidimensional then $ht J \leq n - \text{rank } M$.

Mathematics Subject Classification (1991): 13C15, 13H10, 13A30

Here by the rank of a module M we mean

$$\text{rank } M = \min\{\mu(M_p) \mid p \text{ is a minimal prime of } R\}.$$

Thus, for example, if R is a domain, then this is the usual (torsionfree) rank.

Note that the result is sharp, as can be seen by considering modules M with $\text{projdim } M \leq 1$. Such a module has a minimal presentation

$$0 \longrightarrow R^{n-\text{rank } M} \xrightarrow{\phi} R^n \longrightarrow M \longrightarrow 0,$$

and one may construct examples with the property that a row of ϕ form part of a system of parameters of R ; hence $ht J = \mu(J) = n - \text{rank } M$. (The fact that $\mu(M) - \text{rank } M \leq \dim R$, in the context of Theorem 1, can be seen using [7]; cf. also the proof of Lemma 2 below.)

While this paper was being written, we learned that similar results have been shown by Kwieciński [10] and very recently by Eisenbud, Huneke and Ulrich. In our situation, Kwieciński gives the result that $ht I_1(\phi) \leq n(n - \text{rank } M)$, which is an immediate consequence of the Theorem. He proves this in case R is a regular domain which is finitely generated over a field of characteristic zero; as one might expect from the additional hypotheses, the method of proof is quite different from the current one, which makes use of a deformation argument of [9, proof of Lemma 3.2]. In an early version, we stated the result only in the case for ideals, but the general case is no different.

Before we turn to the proof of Theorem 1, we need a basic fact assuring that the symmetric algebra has the expected dimension.

LEMMA 2. Let R be a noetherian local ring and let M be a finitely generated R -module. If $S(M)$ is equidimensional then $\dim S(M) = \dim R + \text{rank } M$.

Proof. By [7], for any minimal prime p of R , there exists a (unique) minimal prime $T(p)$ of $S(M)$ lying over p , and has dimension

$$\dim S(M)/T(p) = \mu(M_p) + \dim R/p.$$

But as $S(M)$ is equidimensional we must have

$$\dim S(M) = \mu(M_p) + \dim R/p$$

for any minimal prime p of R . It suffices to show that the minimum of $\mu(M_p)$ occurs at a prime with $\dim R/p = \dim R$. But this is clear as the sum $\mu(M_p) + \dim R/p$ is independent of the minimal prime p .

Proof of Theorem 1.

We may assume that J is the ideal generated by the last row of ϕ .

Write

$$S = S(M) = R[T_1, \dots, T_n]/(\ell_1, \dots, \ell_m),$$

where

$$[\ell_1, \dots, \ell_m] = [T_1, \dots, T_n] \cdot \phi,$$

and let $\underline{x} = x_1, \dots, x_{n-1}$ be the images of T_1, \dots, T_{n-1} in S . Set $P = (mS, \underline{x}) \in \text{Spec}(S(M))$, where m denotes the maximal ideal of R . Then there are isomorphisms

$$\begin{aligned} S_P/(\underline{x})S_P &\cong (R[T]/T \cdot JR[T])_{(m)} \\ &\cong (R[T]/JR[T])_{(m)} \\ &\cong (R/J)(T), \end{aligned}$$

where $T = T_n$. Now supposing that $ht J > n - \text{rank } M$, as $S(M)$ is equidimensional and catenary, we have that

$$\begin{aligned} \dim S_P &= \dim S_P/(\underline{x})S_P + ht (\underline{x})S_P \\ &\leq \dim S_P/(\underline{x})S_P + n - 1 \\ &= \dim (R/J)(T) + n - 1 \\ &= \dim R/J + n - 1 \\ &\leq \dim R - ht J + n - 1 \\ &< \dim R - (n - \text{rank } M) + n - 1 \\ &= \dim R + \text{rank } M - 1 \\ &= \dim S - 1 \\ &= \dim S - \dim S/P \\ &= \dim S_P. \end{aligned}$$

This contradiction shows that $ht J \leq n - \text{rank } M$.

THEOREM 3. With the notation as in Theorem 1, assume that $S(M)$ is Cohen-Macaulay and that $ht J \geq n - \text{rank } M$. Then R/J is Cohen-Macaulay.

Proof. Theorem 1 implies that $ht J = n - \text{rank } M$. But now the string of inequalities in the proof are all equalities. Thus $ht (\underline{x})S_P = n - 1$. Since S_P is Cohen-Macaulay, it

follows that \underline{x} is an S_P -sequence. Hence $(R/J)(T) \cong S_P/(\underline{x})S_P$ is Cohen-Macaulay, and thus, by faithfully flat descent, so is R/J .

We note that Theorem 3 also holds if one replaces “Cohen-Macaulay” by “Gorenstein”. A similar remark holds for “complete intersection”, but this is a less interesting condition for the symmetric algebra, as it means, at least if R is regular, that $\text{projdim } M \leq 1$ (cf. [13]). In this case the result would be clear: trivially $\mu(J) \leq \#$ columns of $\phi = n - \text{rank } M$, thus if $ht J \geq n - \text{rank } M$, then $\mu(J) = ht J$, hence J and R/J are complete intersections.

We make one final note about the proof. This is the observation that we did not need the full strength that $S(M)$ is Cohen-Macaulay, but that a certain localization has this property. We will make use of this remark in a moment.

Applications to Residual Intersections

In the case of ideals, the previous results can be conveniently described using the notion of residual intersection. We find it useful to give the translation into this language.

Recall that a proper ideal J is said to be an s -residual intersection of I if there are elements a_1, \dots, a_s in I such that $J = (a_1, \dots, a_s) : I$ and $ht J \geq s$. If in addition $ht I + J > s$ then J is called a geometric s -residual intersection. (To avoid trivial cases, one usually also assumes that $s \geq ht I$, but we will find it convenient here not to impose this convention.) To study upper bounds on the heights of colon ideals one may often restrict attention to residual intersections. Actually, one would even like to know when an s -residual intersection has the “expected height” s , and when it is Cohen-Macaulay [6],[5],[8],[11].

We set $A = (a_1, \dots, a_s)$ and consider residual intersections $J = A : I$ with I/A cyclic. Such a J may obviously be chosen as a row of a matrix presenting I . Hence Theorems 1 and 3 now yield

COROLLARY 4. Let R be a universally catenary local ring, let I be an R -ideal with $ht I > 0$, and let $J = A : I$ be an s -residual intersection of I with I/A cyclic.

If $S(I)$ is equidimensional then $ht J = s$; if in addition $\text{Proj } (S(I))$ is Cohen-Macaulay then so is R/J .

Of course both conditions in the corollary are satisfied whenever $S(I)$ is Cohen-Macaulay. (How much weaker the conditions in the corollary are, however, is not so clear to the author.)

As an application, we now wish to give a residual intersection characterization of the Cohen-Macaulayness of the Rees algebra, for ideals of linear type. This may be viewed as a partial generalization of a result [3, Corollary 6.4] on d -sequences.

Recall that an ideal I of a noetherian ring is said to be of linear type if the canonical epimorphism $S(I) \rightarrow \mathcal{R}(I)$ to the Rees algebra $\mathcal{R}(I) = \bigoplus_{i \geq 0} I^i \cong R[It]$ is an isomorphism; in particular, this implies an isomorphism $S_{R/I}(I/I^2) \cong gr_I(R) = \bigoplus_{i \geq 0} I^i/I^{i+1}$. For an ideal of linear type and positive height in a local Cohen-Macaulay ring, it is known that the Cohen-Macaulay property of $\mathcal{R}(I)$ is equivalent to that of $gr_I(R)$ (cf. e.g. [1]). Thus we may concentrate on the Cohen-Macaulayness of $gr_I(R)$.

We will exploit a basic example of [6], 4.3, which shows that, in the case of an ideal of linear type in a local Cohen-Macaulay ring, the extended Rees algebra $R[It, t^{-1}]$ is defined by an ideal J in $R[T_1, \dots, T_n, U]$ which is a residual intersection of (I, U) . More precisely, if $n = \mu(I)$, then one has $J = A : (I, U)$, where J is a geometric n -residual intersection and $(I, U)/A$ is cyclic.

THEOREM 5. Let R be a local Cohen-Macaulay ring and let I be an R -ideal of linear type. Then the following are equivalent.

- (a) $gr_I(R)$ is Cohen-Macaulay ;
- (b) for any local Cohen-Macaulay faithfully flat extension R' of R , and any $x \in R'$ regular on $gr_{IR'}(R')$, every residual intersection $J = A : (I, x)$ of (I, x) in R' , with $(I, x)/A$ cyclic, is Cohen-Macaulay.

Proof. If (b) holds then in particular taking $R' = R[T_1, \dots, T_n, U]_{(m_R, T_1, \dots, T_n, U)}$, and $x = U$, as we have said above, we have that $R[It, t^{-1}] \cong R[T_1, \dots, T_n, U]/J$ is Cohen-Macaulay (as it is Cohen-Macaulay locally at its unique graded maximal ideal). Hence $gr_I(R) \cong R[It, t^{-1}]/(t^{-1})$ is Cohen-Macaulay.

Conversely, suppose that $gr_I(R)$ is Cohen-Macaulay. We claim that $gr_{(I, x)}(R')$ is Cohen-Macaulay. First, note that by the flatness of the morphism $R \rightarrow R'$ we have that IR' is still of linear type. Further, since $R \rightarrow T$ is a faithfully flat map of Cohen-Macaulay rings, it follows that by base change (after localizing at the irrelevant maximal ideals) that $gr_I(R) \rightarrow gr_{IR'}(R')$ has the same property; in particular, $gr_{IR'}(R')$ is Cohen-Macaulay. Now (I, x) is still of linear type [12, Proposition 2.5], and $gr_{(I, x)}(R') \cong gr_{IR'}(R')/(x)gr_{IR'}(R')$. Hence $gr_{(I, x)}(R')$ is Cohen-Macaulay, proving the claim. Thus we may conclude that $\mathcal{R}((I, x))$ is Cohen-Macaulay. Now we

may apply Corollary 4 to $(I, x) \subset R'$ as $S_{R'}((I, x)) \cong \mathcal{R}((I, x))$ is Cohen-Macaulay, to obtain the required result.

We next consider the Gorenstein case. This partially generalizes [4, Theorem 1.3].

THEOREM 6. Let R be a local Gorenstein ring and let I be an R -ideal of linear type. Then the following are equivalent.

- (a) $gr_I(R)$ is Gorenstein ;
- (b) for any local Cohen-Macaulay faithfully flat extension R' of R , and any $x \in R'$ regular on $gr_{IR'}(R')$, every residual intersection $J = A : (I, x)$ of (I, x) in R' , with $(I, x)/A$ cyclic, is Gorenstein.

Proof. If (b) holds then $R[It, t^{-1}] \cong R[T_1, \dots, T_n, U]/J$ is Gorenstein, hence so is $gr_I(R) \cong R[It, t^{-1}]/(t^{-1})$.

Conversely, suppose that $gr_I(R)$ is Gorenstein. Then as in the proof of Theorem 5 it follows that $gr_{(I, x)}(R')$ is Gorenstein. Then, as is well-known, the blow-up $\text{Proj}(\mathcal{R}((I, x)))$ is Gorenstein. Since (I, x) is of linear type, the result now follows from the Gorenstein version of Corollary 4.

We now wish to give a partial extension of Corollary 4. As an s -residual intersection in general can have height greater than s , obviously one can not simply delete the equidimensional assumption on the symmetric algebra. However, it turns out that if we know already that the residual intersection has height s , and is unmixed (and geometric as well), then we may replace the assumption that $S(I)$ is Cohen-Macaulay with the Cohen-Macaulayness of the Rees algebra. This is actually a partial generalization, because under these extra hypotheses, the Cohen-Macaulayness of the symmetric algebra would already imply that it is the Rees algebra ([2], 6.8).

PROPOSITION 7. Let R be a local Cohen-Macaulay ring, let I be an R -ideal with $ht I > 0$, and let $J = A : I$ be a geometric s -residual intersection of I with I/A cyclic, and assume that J is unmixed and that $ht J = s$.

If $\text{Proj}(\mathcal{R}(I))$ is Cohen-Macaulay then so is R/J .

Proof. Write $A = (f_1, \dots, f_s)$, $I = (f_1, \dots, f_{s+1})$ and choose a presentation

$$\mathcal{R}(I) \cong R[T_1, \dots, T_{s+1}]/Q,$$

where Q is the ideal generated by all forms $F(T_1, \dots, T_{s+1})$ vanishing on f_1, \dots, f_{s+1} .

The proof will follow that of Theorem 1, once we show the claim that

$$Q \subset (T_1, \dots, T_s, T_{s+1}J).$$

Indeed, as $T_{s+1}J$ is contained in Q modulo (T_1, \dots, T_s) , this containment would imply that $(Q, T_1, \dots, T_s) = (T_1, \dots, T_s, T_{s+1}J)$, giving the usual isomorphism locally at P , modulo the sequence \underline{x} .

To verify the claim, let $F \in Q$ be a form of degree $k \geq 1$. Then modulo (T_1, \dots, T_s) we may write $F \equiv aT_{s+1}^k$ with $a \in R$. We must show that $a \in J$. By evaluating, we see that $af_{s+1}^k \in (f_1, \dots, f_s)$, hence $af_{s+1}^{k-1} \in (f_1, \dots, f_s) : (f_{s+1}) = J$. Now if $a \notin J$, then $a \notin q$ for some p -primary component q of J . But then as $af_{s+1}^{k-1} \in J \subset q$, in particular p would contain f_{s+1} , hence $I + J$. Thus, as J is geometric, $ht p \geq ht I + J > s$, which contradicts the unmixedness of J . This proves the claim.

Notice that (as in the original argument [9], 3.2) as I has positive height, \mathcal{R} is equidimensional and $\dim \mathcal{R} = \dim R + 1$. Furthermore, $P = (m, \underline{x}) \in \text{Proj}(\mathcal{R}(I))$, hence in particular the local ring S_P is Cohen-Macaulay. Now the result follows exactly as in the proof of Theorems 1 and 3.

The assumption that the blow-up $\text{Proj}(\mathcal{R}(I))$ is Cohen-Macaulay holds of course if $\mathcal{R}(I)$ is Cohen-Macaulay; more generally, it is well-known that the blow-up is Cohen-Macaulay if the associated graded ring $gr_I(R)$ is Cohen-Macaulay.

We should point out again that Theorem 5 (as well as Corollary 4) holds if we replace the condition "Cohen-Macaulay" by "Gorenstein" or "complete intersection." One should note, however, that it is false for "regular"; this is due to the fact that regularity is not preserved under specialization. If the blow-up is smooth though, in some cases one can show that a *sufficiently general* residual intersection is regular ([9], 4.1).

We should also say that, of course, that the proposition does not hold without the assumption that J is unmixed, that is to say that even if $\mathcal{R}(I)$ is Cohen-Macaulay, a geometric s -residual intersection of height s need not be Cohen-Macaulay. One way, for instance, to construct such examples is to use again the result [6], 4.3, which (more generally without the assumption of linear type) gives the so-called extended symmetric algebra as a geometric n -residual intersection of height n .

References

- [1] I. M. Aberbach, C. Huneke and N. V. Trung, Reduction numbers, Briançon-Skoda theorems and the depth of Rees rings, *Compositio Math.* **97**(1995), 403-434.
- [2] J. Herzog, A. Simis and W. V. Vasconcelos, Approximation complexes of blowing-up rings II, *J. Algebra* **82**(1983), 53-83.
- [3] J. Herzog, A. Simis and W. V. Vasconcelos, On the arithmetic and homology of algebras of linear type, *Trans. Amer. Math. Soc.* **283** (1984), 661-683.
- [4] J. Herzog, A. Simis and W. V. Vasconcelos, On the canonical module of the Rees algebra and the associated gradedring of an ideal, *J. Algebra* **105**(1987), 285-302.
- [5] J. Herzog, W. V. Vasconcelos and R. Villarreal, Ideals with sliding depth, *Nagoya Math. J.* **99**(1985), 159-172.
- [6] C. Huneke, Strongly Cohen-Macaulay schemes and residual intersections, *Trans. Amer. Math. Soc.* **277**(1983), 739-763.
- [7] C. Huneke and M. Rossi, The dimension and components of symmetric algebras, *J. Algebra* **98**(1986), 200-210.
- [8] C. Huneke and B. Ulrich, Residual intersections, *J. Reine Angew. Math.* **390**(1988), 1-20.
- [9] C. Huneke, B. Ulrich and W. V. Vasconcelos, On the structure of certain normal ideals, *Compositio Math.* **84** (1992), 25-42.
- [10] M. Kwieciński, Bounds for codimensions of Fitting ideals, *J. Algebra* **194**(1997), 378-382.
- [11] B. Ulrich, Artin-Nagata properties and reductions of ideals, *Commutative algebra: syzygies, multiplicities, and birational algebra (South Hadley, MA, 1992)*, 373-400, Contemp. Math., 159, Amer. Math. Soc., Providence, RI, 1994.
- [12] G. Valla, On the symmetric and Rees algebras of an ideal, *Manuscripta Math.* **30** (1980), 239-255.
- [13] W. V. Vasconcelos, Symmetric Algebras, *Commutative algebra (Salvador, 1988)*, 115-160, Lecture Notes in Math., 1430, Springer-Verlag, New York, 1990.

Department of Mathematical Sciences, University of Arkansas, Fayetteville, AR 72701
mark@math.uark.edu